



Antiapoptotic Effects of Continuous Training and Selenium Consumption on the Liver Tissue of Cadmium-Exposed Rats

Hossein Humayun Fard¹, Seyed Ali Hosseini^{2,*}, Mohammad Ali Azarbayjani³ and Masoud Nikbakht⁴

¹Department of Sport Physiology, Shoushtar Branch, Islamic Azad University, Shoushtar, Iran

²Department of Sport Physiology, Marvdasht Branch, Islamic Azad University, Marvdasht, Iran

³Department of Sport Physiology, Central Tehran Branch, Islamic Azad University, Tehran, Iran

⁴Department of Exercise Physiology, Faculty of Sport Sciences, Shahid Chamran University of Ahvaz, Ahvaz, Iran

*Corresponding author: Department of Sport Physiology, Marvdasht Branch, Islamic Azad University, Marvdasht, Iran. Email: alihoseini_57@miau.ac.ir

Received 2019 March 05; Revised 2019 March 17; Accepted 2019 April 17.

Abstract

Background: Cadmium has negative effects on various tissues of the body while selenium has antioxidant activities. Moreover, the benefits of exercise training for apoptotic factors including caspase-3 and cytochrome c, as well as cyclin D, have been regarded.

Objectives: Therefore, the present study aimed to evaluate the anti-apoptotic effects of continuous training and selenium consumption on the liver tissue of cadmium-exposed rats.

Methods: In this study, 25 rats were selected and randomly divided into five groups of five rats including: (1) control, (2) selenium consumption, (3) continuous training, (4) continuous training with selenium consumption, and (5) sham. For eight weeks, groups 1 to 4 received peritoneal cadmium (2 mg/kg) daily; groups 2 and 4 received peritoneal selenium (0.23 mg/kg) daily; and groups 3 and 4 performed continuous training three sessions per week. Caspase-3, cytochrome c, and cyclin D were measured at the protein level.

Results: Cadmium consumption significantly increased the protein levels of caspase-3 and cytochrome c and decreased cyclin D in rats ($P = 0.001$). Selenium consumption and continuous training significantly decreased the protein levels of caspase-3 and cytochrome c and increased cyclin D in rats exposed to cadmium ($P = 0.001$). Continuous training along with selenium consumption had interactive effects on increasing the protein levels of cyclin D in rats exposed to cadmium ($P = 0.02$).

Conclusions: It appears that continuous training and selenium consumption have interactive anti-apoptotic effects in the liver tissue of cadmium-exposed rats.

Keywords: Liver, Apoptosis, Continues Training, Selenium, Cadmium

1. Background

Cadmium is one of the environmental pollutants that produces high levels of toxicity in the liver, thus increasing liver enzymes and inducing apoptosis from the pathway of caspase activation (1). During apoptosis, due to the fragmentation and detachment of the protein from the small and large subunits, procaspase is converted into an active caspase. The active caspase affects other specific sites of caspases, which can activate their lower caspases, causing a proteolytic cascade. Next, the active caspases break other key proteins in the cell. Another group of caspases can break proteins that inhibit DNA degrading enzymes; thus, DNA breaks down in the nucleus of the cell with the release of DNAases (2).

Cadmium damages many tissues, such as the pancreas, liver, lungs, testicles, and kidneys. The main tissue of the body that is affected by cadmium is the liver because this

metal increases oxidative pressure (3). A study has shown that in rats exposed to dextran sodium sulfate, this toxic substance increased the oxidative stress and apoptotic-inducing factors. In contrast, in rats that performed continuous swimming one to 1.5 hours daily for seven weeks, apoptotic factors such as caspase-3, cytochrome c, and Bax showed significantly lower values than in the group receiving dextran sodium sulfate (4). In the same vein, it was reported that the Bax gene expression and the ratio of Bax to Bcl2, which are factors in the production of caspase-3, were lower in rats that performed running on a treadmill for 12 weeks than in those that did not perform any physical activity (5).

Selenium, as an essential ingredient, is naturally occurring in soil, water, and many foods, and given its antioxidant properties, it is essential for health (6). According to the literature, cadmium consumption increases caspase-

3 and cytochrome c from the pathway of peroxisome proliferator-activated receptor gamma/phosphoinositide 3-kinase/Akt (PPAR- γ /PI3K/Akt) and thus, induces apoptosis in the pancreatic tissue; however, in groups receiving selenium, the effects of apoptosis induced by cadmium had a significant decrease (7). In the review of the effects of selenium and cadmium on apoptosis, researchers concluded that selenium is effective in reducing the induction of cadmium-induced apoptosis (8). In mammalian cells, there are three types of cyclins, D1, D2, and D3 (9). In a study of embryonic fibroblasts cells, it was concluded that cadmium increases cyclin levels (10). Moreover, in another study conducted on rats performing endurance swimming, exercise training reduced the apoptosis-inducing factors in the liver tissue of cadmium-receiving rats (11).

2. Objectives

Considering the contradictions reported in some studies, as well as the effects of selenium on reducing the production of free radicals and inflammation and the strengthening of antioxidant defense devices, and the effects of continuous training on reducing the apoptosis-inducing factors, the aim of this study was to determine the interactive effects of continuous exercise and selenium consumption on caspase-3, cytochrome c, and cyclin D in the liver tissue of cadmium-exposed rats.

3. Methods

The present experimental study was conducted at the Animal Lab of Marvdasht Branch of Islamic Azad University, in which 25 male rats (Sprague Dawley) were purchased from the Animal Breeding Site of the university. In order to ensure that the rats were adapted to the laboratory environment, they were kept in clear polycarbonate cages with humidity of 45% to 55%, the dark/light cycle of 12/12 hours, and temperature of $23^{\circ}\text{C} \pm 2^{\circ}\text{C}$. During the study, a standard diet of pellets and water was given to rats *ad libitum*. After one week, rats were randomly divided into five groups of five rats, including: (1) control, (2) selenium, (3) continuous training, (4) continuous training with selenium consumption, and (5) sham. The study was approved by the Local Ethics Committee of the Sport Sciences Research Institute of Iran.

3.1. Cadmium and Selenium Consumption

Groups 1 to 4 received cadmium (2 mg/kg) daily for two weeks peritoneally (12). A cadmium solution was prepared by dissolving cadmium chloride in normal saline on a daily basis to administer the aforementioned dosage.

During the study, rats in the sham group received 2 mg/kg of cadmium-free solution (normal saline) daily. Groups 2 and 4 received peritoneal selenium (0.23 mg/kg) in the form of sodium selenite daily (13). Groups 3 and 4 performed continuous training on a rat's treadmill three sessions a week for eight weeks (14).

3.2. Continuous Training Protocol

To evaluate the maximum speed running, we used Bedford et al. incremental test (15) as standardized by Leandro et al. (16) for rats. The continuous training protocol consisted of three stages of warming up (5 minutes), the main training, and cooling down (5 minutes). Rats in the continuous training groups first warmed up on the treadmill for 5 minutes at an intensity of 50% - 60% of $\text{VO}_{2\text{max}}$. Then, they performed continuous training at an intensity of 65% of $\text{VO}_{2\text{max}}$ in the first week, 70% of $\text{VO}_{2\text{max}}$ in the second week, and 75% of $\text{VO}_{2\text{max}}$ since the third week onward. The duration of the continuous training was 6 minutes in the first week, 14 minutes in the second week, 22 minutes in the third week, and 30 minutes from the fourth week to the eighth week. At the end of each session, 5 minutes of cooling down at an intensity of 50% to 60% of $\text{VO}_{2\text{max}}$ was performed. Finally, 48 hours after the last training session and last injection of cadmium and selenium in the eighth week, the rats were anesthetized by peritoneal injection of ketamine and xylazine and their liver tissues were removed to measure the variables of research.

3.3. Measuring the Research Variables

In order to measure caspase-3, cytochrome c, and cyclin D by Enzyme-linked immunosorbent assay (ELISA), the liver tissue was placed in a microtube after washing with normal saline and freezing in a nitrogen tank at -80°C . In accordance with the instructions of the kit manufacturer, 100 mg of the frozen tissue was ground into a powder to measure the protein level of caspase-3 (CSB-E08857r. Cusabio Biotech Co., USA) and cytochrome c (CSB-E14281r. Cusabio Biotech Co., USA). For the homogenization of tissues, Phosphate-buffered Saline (PBS) solution was used to combine one milliliter PBS and one mg of the tissue. After storing the mixture at -20°C , freezing, and melting twice, it was centrifuged for 5 minutes at a temperature of $2^{\circ}\text{C} - 8^{\circ}\text{C}$ at a speed of 5000 rpm to break the cell wall. Then, cytochrome c and caspase-3 were measured immediately using standard solutions. To measure the protein level of cyclin D, the ELISA kit (SEA585Ra 96 Tests, Cloud-Clone Corp., USA) was used according to the manufacturer's instructions.

3.4. Statistical Analysis

The Shapiro-Wilk test was used to investigate the normal distribution of data. To determine the effect of continuous training and selenium consumption on research variables, the two-way ANOVA and independent samples *t*-test were used in SPSS (version 21) software. The significance level was set at $P \leq 0.05$.

4. Results

The levels of cytochrome c, cyclin D, and caspase-3 in the five study groups are presented in Figures 1 to 3. Table 1 shows that cytochrome c and caspase-3 were significantly higher in the control group than in the sham group ($P = 0.001$). Nevertheless, cyclin D was significantly lower in the control group than in the sham group ($P = 0.001$). Therefore, cadmium consumption significantly increased the protein levels of caspase-3 and cytochrome c and decreased cyclin D in rats.




Figure 1. Levels of caspase-3 in the five study groups




Figure 2. Levels of cytochrome c in the five study groups




Figure 3. Levels of cyclin D in the five study groups

Table 1. Results of Independent Samples *t*-Test to Compare Caspase-3, Cytochrome c, and Cyclin D in Control and Sham Groups

Variable	<i>t</i>	df	P Value
Caspase-3	-5.89	8	0.001
Cyclin D	26.31	8	0.001
Cytochrome c	-18.12	8	0.001

Table 2 shows that selenium ($P = 0.001$) and continuous training ($P = 0.001$) had significant effects on decreasing the protein levels of caspase-3 in rats exposed to cadmium. However, the interaction of continuous training and selenium was not significant ($P = 0.12$).

Selenium ($P = 0.001$) and continuous training ($P = 0.01$) had significant effects on increasing the protein level of cyclin D in rats exposed to cadmium. Moreover, the interaction of continuous training and selenium was significant ($P = 0.02$).

In addition, selenium ($P = 0.001$) and continuous training ($P = 0.001$) had significant effects on decreasing the protein levels of cytochrome c in rats exposed to cadmium. However, the interaction of continuous training and selenium was not significant ($P = 0.14$).

5. Discussion

The findings of the present study revealed that cadmium consumption significantly increased cytochrome c and caspase-3 apoptotic factors and decreased cyclin D as a factor promoting cell division in rats. Cadmium is a pervasive environmental pollutant and is one of the most important toxins of the liver. By activating the pathway of nuclear factor erythroid 2-related factor 2 (Nrf2), this toxic substance could cause caspase-dependent apoptosis and liver damage in cadmium-receiving rats (1), which is parallel with the findings of the present study. Based on the

Table 2. Results of Two-Way ANOVA on the Effects of Continuous Training with Selenium Consumption on Cytochrome c, Caspase-3, and Cyclin D in Rats Exposed to Cadmium

Source	Sum of Squares	df	Mean Square	F	P Value	Partial Eta Squared
Caspase-3						
Selenium consumption	9.59	1	9.59	36.02	0.001	0.69
Training	4.69	1	4.69	17.65	0.001	0.52
Interaction of training and selenium consumption	0.699	1	0.69	2.62	0.12	0.14
Cyclin D						
Selenium consumption	19.99	1	19.99	354.34	0.001	0.95
Training	0.40	1	0.40	7.22	0.01	0.31
Interaction of training and selenium consumption	0.36	1	0.36	6.48	0.02	0.28
Cytochrome c						
Selenium consumption	22.37	1	22.37	286.61	0.001	0.94
Training	2.32	1	2.32	29.77	0.001	0.65
Interaction of training and selenium consumption	0.18	1	0.18	2.33	0.14	0.12

study of cadmium-exposed liver cells, cadmium increases caspase-3 and cytochrome c, and induces apoptosis (17), which are parallel with the findings of the present study.

Apoptosis is one of the most common ways to remove unwanted cells, which occurs in the body of multicellular and even monocellular organisms. In this process, the cell is responsible for its own death; thereby, it is also called cell suicide. By applying this process, the immune system performs many of its anti-antigen activities. Caspases are the most active executive members of the set. The name of caspase (cysteine aspartate-protease) is taken from their functions. According to a theory, the Apaf-1, caspase-9, and cytochrome c complexes can activate caspase-9 activation. Caspase-9 enzyme activation results in caspase-3 activity. Caspase-3 performs various actions as the executive member of this set (18).

Cyclin D is a promoter of the cell cycle during the G1 stage, which drives the cell cycle into the S stage. This protein plays a key role in cell proliferation, cell growth regulation, and mitochondrial activity fluctuations (19). Based on research conducted on prostatic epithelial cells, cadmium reduces the protein level of cyclin D (20), which is parallel with the findings of the present study.

The current study also showed that continuous training significantly decreased caspase-3 and cytochrome c and increased cyclin D in rats exposed to cadmium. In this regard, swimming endurance training in pregnant rats resulted in a significant reduction in apoptotic indices in the liver tissue of cadmium-receiving neonatal rats (21), which is parallel with the findings of the present study.

Moreover, in line with the findings of the present study, it has been shown that continuous training increases Bcl2 as an anti-apoptotic factor and decreases Bax as an inducer

of apoptosis in cadmium-exposed rats (22). In addition, some studies have concluded that physical exercise may increase cyclin D levels in nerve cells of rats (23). In this study, selenium consumption resulted in a significant reduction in caspase-3 and cytochrome c and a significant increase in cyclin D in cadmium-exposed rats.

Selenium is an essential ingredient in selenocysteines. The selenoproteins, including glutathione peroxidase (GPXs), effectively perform antioxidant activities and protect against the reactive oxygen and nitrogen species (ROS and RNS); however, over-consumption of selenium can also be toxic to the body (24). In many studies, cell proliferation, control, or death and hemostasis in many tissues, as well as the control of cell cycle and the process of apoptosis, have been reported to be the effects of selenium in the body (25).

Selenium deficiency increases the expression of miR-193b-3p (microRNA-193b-3p) gene, which reduces the gene expression of Mastermind-like protein 1 (MAML1) and thus increases hepatocyte apoptosis (26). In Binte Hossain et al.'s study, the levels of glutathione and glutathione peroxidase 1 (GPx1) were higher in groups co-administered with cadmium and selenium than in the cadmium-only group. Selenium also reduced cytochrome c and decreased apoptosis (27).

Concerning the interactive effects, in this study, continuous training along with selenium consumption had interactive effects on the increase of cyclin D in rats exposed to cadmium; however, there was no interactive effect on the reduction of caspase-3 and cytochrome c. Even though studies in the literature have investigated the effect of continuous training or selenium consumption alone on the levels of apoptotic factors of caspase-3, cytochrome c,

and cyclin D in various organisms, including cadmium-exposed rats, no study has evaluated the interactional effects of selenium with continuous exercise on caspase-3, cytochrome c, and cyclin D in cadmium-exposed rats, making it impossible to compare the results. Therefore, further studies are needed to be fully aware of this interactive effect.

There are some limitations to this study, such as the lack of controlling for calories in the tissues of the rats, the multiplicity of the factors signaling the apoptotic pathway, and cell proliferation. Therefore, it is suggested that future studies investigate the different doses of cadmium and selenium in different tissues and examine other factors of apoptosis and cell proliferation to achieve more reliable and dependable results. In addition, the effect of cadmium via peritoneal, oral, and inhalational administrations should be compared.

5.1. Conclusions

Based on the findings of the current study, continuous training and selenium consumption alone can decrease the apoptosis protein factors of caspase-3 and cytochrome c and increase cyclin D in the liver tissue of rats exposed to cadmium. In addition, continuous training and selenium consumption have interactive effects on cyclin D in cadmium-exposed rats; however, the interactive effects were not significant in reducing caspase-3 and cytochrome c.

Acknowledgments

The present study is part of a doctoral dissertation approved by Shoushtar Islamic Azad University. The authors express their gratitude and appreciation to the Research and Technology Department of the University.

Footnotes

Authors' Contribution: All authors equally contributed to the writing and revision of this paper.

Conflict of Interests: The authors declare that they have no conflict of interests.

Ethical Approval: The study received approval from the Sports Sciences Research Institute of Iran with ethics code IR.SSRI.REC.1397.38.

Funding/Support: Shoushtar Islamic Azad University funded the study.

References

- Gong ZG, Wang XY, Wang JH, Fan RF, Wang L. Trehalose prevents cadmium-induced hepatotoxicity by blocking Nrf2 pathway, restoring autophagy and inhibiting apoptosis. *J Inorg Biochem.* 2019;**192**:62-71. doi: [10.1016/j.jinorgbio.2018.12.008](https://doi.org/10.1016/j.jinorgbio.2018.12.008). [PubMed: [30599315](https://pubmed.ncbi.nlm.nih.gov/30599315/)].
- Ghadirian S, Naji T. [Apoptosis and Flow cytometry]. *Tashkhis Aze-mayeshgahi.* 2018;**13**:33-0. Persian.
- Arroyo VS, Flores KM, Ortiz LB, Gomez-Quiroz LE, Gutierrez-Ruiz MC. Liver and cadmium toxicity. *Drug Metab Toxicol.* 2012;**2**:2-7.
- Qin L, Yao ZQ, Chang Q, Zhao YL, Liu NN, Zhu XS, et al. Swimming attenuates inflammation, oxidative stress, and apoptosis in a rat model of dextran sulfate sodium-induced chronic colitis. *Oncotarget.* 2016;**8**(5). doi: [10.18632/oncotarget.14080](https://doi.org/10.18632/oncotarget.14080).
- Jafari A, Pourrazi H, Nikookheslat S, Baradaran B. Effect of exercise training on Bcl-2 and bax gene expression in the rat heart. *Gene Cell Tissue.* 2015;**2**(4). doi: [10.17795/gct-32833](https://doi.org/10.17795/gct-32833).
- Wang N, Tan HY, Li S, Xu Y, Guo W, Feng Y. Supplementation of micronutrient selenium in metabolic diseases: Its role as an antioxidant. *Oxid Med Cell Longev.* 2017;**2017**:7478523. doi: [10.1155/2017/7478523](https://doi.org/10.1155/2017/7478523). [PubMed: [29441149](https://pubmed.ncbi.nlm.nih.gov/29441149/)]. [PubMed Central: [PMC5758946](https://pubmed.ncbi.nlm.nih.gov/PMC5758946/)].
- Jin X, Jia T, Liu R, Xu S. The antagonistic effect of selenium on cadmium-induced apoptosis via PPAR-gamma/P13K/Akt pathway in chicken pancreas. *J Hazard Mater.* 2018;**357**:355-62. doi: [10.1016/j.jhazmat.2018.06.003](https://doi.org/10.1016/j.jhazmat.2018.06.003). [PubMed: [29909169](https://pubmed.ncbi.nlm.nih.gov/29909169/)].
- Wan N, Xu Z, Liu T, Min Y, Li S. Ameliorative effects of selenium on cadmium-induced injury in the chicken ovary: Mechanisms of oxidative stress and endoplasmic reticulum stress in cadmium-induced apoptosis. *Biol Trace Elem Res.* 2018;**184**(2):463-73. doi: [10.1007/s12011-017-1193-x](https://doi.org/10.1007/s12011-017-1193-x). [PubMed: [29090375](https://pubmed.ncbi.nlm.nih.gov/29090375/)].
- Qie S, Diehl JA. Cyclin D1, cancer progression, and opportunities in cancer treatment. *J Mol Med (Berl).* 2016;**94**(12):1313-26. doi: [10.1007/s00109-016-1475-3](https://doi.org/10.1007/s00109-016-1475-3). [PubMed: [27695879](https://pubmed.ncbi.nlm.nih.gov/27695879/)]. [PubMed Central: [PMC5145738](https://pubmed.ncbi.nlm.nih.gov/PMC5145738/)].
- Xie H, Wang J, Jiang L, Geng C, Li Q, Mei D, et al. ROS-dependent HMG2 upregulation mediates Cd-induced proliferation in MRC-5 cells. *Toxicol In Vitro.* 2016;**34**:146-52. doi: [10.1016/j.tiv.2016.04.001](https://doi.org/10.1016/j.tiv.2016.04.001). [PubMed: [27071802](https://pubmed.ncbi.nlm.nih.gov/27071802/)].
- Mirdar Harijani S, Musavi NGR, Hamidian G, Asgharzadeh Oliaei HA. [Effect of endurance swimming training and silymarin treatment on changes in liver apoptotic index in pregnant rats exposed to cadmium]. *J Rafsanjan Univ Med Sci.* 2014;**13**(8):705-14. Persian.
- Rahimi Kazerouni M, Mokhtari M, Shariati M, Rahimi Kazerouni S. [The effect of *Oliveria decumbens* aqueous extract on functional factors and tissue changes in adult male rats under cadmium chloride poisoning]. *J Physiol Animal Evolut.* 2015;**29**(2):42-33. Persian.
- Dzobo K, Naik YS. Effect of selenium on cadmium-induced oxidative stress and esterase activity in rat organs. *South Africa J Sci.* 2013;**109**(5/6):1-8. doi: [10.1590/sajs.2013/965](https://doi.org/10.1590/sajs.2013/965).
- Fathi E, Noor Shahi M, Haghparast A, Falah Hosseini H. [Effect of eight-week aerobic continuous and high intensity interval training on levels of SIRT3 in skeletal muscle tissue of Wistar rats]. *Sport Exer Physiol.* 2015;**16**:1277-89. Persian.
- Bedford TG, Tipton CM, Wilson NC, Oppliger RA, Gisolfi CV. Maximum oxygen consumption of rats and its changes with various experimental procedures. *J Appl Physiol Respir Environ Exerc Physiol.* 1979;**47**(6):1278-83. doi: [10.1152/jappl.1979.47.6.1278](https://doi.org/10.1152/jappl.1979.47.6.1278). [PubMed: [536299](https://pubmed.ncbi.nlm.nih.gov/536299/)].
- Leandro CG, Levada AC, Hirabara SM, Manhaes-de-Castro R, De-Castro CB, Curi R, et al. A program of moderate physical training for Wistar rats based on maximal oxygen consumption. *J Strength Cond Res.* 2007;**21**(3):751-6. doi: [10.1519/R-20155.1](https://doi.org/10.1519/R-20155.1). [PubMed: [17685693](https://pubmed.ncbi.nlm.nih.gov/17685693/)].
- Lawal AO, Marnewick JL, Ellis EM. Heme oxygenase-1 attenuates cadmium-induced mitochondrial-caspase 3-dependent apoptosis in human hepatoma cell line. *BMC Pharmacol Toxicol.* 2015;**16**:41. doi: [10.1186/s40360-015-0040-y](https://doi.org/10.1186/s40360-015-0040-y). [PubMed: [26670903](https://pubmed.ncbi.nlm.nih.gov/26670903/)]. [PubMed Central: [PMC4681021](https://pubmed.ncbi.nlm.nih.gov/PMC4681021/)].

18. Honardoost M, Soleimanjahi H, Rajaei F. [Apoptosis: Programmed cell death]. *J Qazvin Univ Med Sci*. 2013;**17**(3):48–57. Persian.
19. Ramos-Garcia P, Gil-Montoya JA, Scully C, Ayen A, Gonzalez-Ruiz L, Navarro-Trivino FJ, et al. An update on the implications of cyclin D1 in oral carcinogenesis. *Oral Dis*. 2017;**23**(7):897–912. doi: [10.1111/odi.12620](https://doi.org/10.1111/odi.12620). [PubMed: [27973759](https://pubmed.ncbi.nlm.nih.gov/27973759/)].
20. Liu Q, Ji X, Ge Z, Diao H, Chang X, Wang L, et al. Role of connexin 43 in cadmium-induced proliferation of human prostate epithelial cells. *J Appl Toxicol*. 2017;**37**(8):933–42. doi: [10.1002/jat.3441](https://doi.org/10.1002/jat.3441). [PubMed: [28176351](https://pubmed.ncbi.nlm.nih.gov/28176351/)].
21. Mirdar Harijani S, Asgharzadeh Oliaei HA, Hamidian G, Musavi N. [The effect of swimming endurance training and silymarin supplementation during pregnancy on maternal cadmium exposure - induced apoptosis in hepatocytes of rat neonates]. *J Pract Stud Biosci Sport*. 2014;**2**(3):9–18. Persian.
22. Ghajari H, Hosseini SA, Farsi S. The effect of endurance training along with cadmium consumption on Bcl-2 and bax gene expressions in heart tissue of rats. *Ann Milit Health Sci Res*. 2019;**17**(1). doi: [10.5812/amh.86795](https://doi.org/10.5812/amh.86795).
23. Liu T, Yang T, Xu Z, Tan S, Pan T, Wan N, et al. MicroRNA-193b-3p regulates hepatocyte apoptosis in selenium-deficient broilers by targeting MAML1. *J Inorg Biochem*. 2018;**186**:235–45. doi: [10.1016/j.jinorgbio.2018.06.013](https://doi.org/10.1016/j.jinorgbio.2018.06.013). [PubMed: [29990747](https://pubmed.ncbi.nlm.nih.gov/29990747/)].
24. Prabhu KS, Lei XG. Selenium. *Adv Nutr*. 2016;**7**(2):415–7. doi: [10.3945/an.115.010785](https://doi.org/10.3945/an.115.010785). [PubMed: [26980826](https://pubmed.ncbi.nlm.nih.gov/26980826/)]. [PubMed Central: [PMC4785479](https://pubmed.ncbi.nlm.nih.gov/PMC4785479/)].
25. Zeng H. Selenium as an essential micronutrient: roles in cell cycle and apoptosis. *Molecules*. 2009;**14**(3):1263–78. doi: [10.3390/molecules14031263](https://doi.org/10.3390/molecules14031263). [PubMed: [19325522](https://pubmed.ncbi.nlm.nih.gov/19325522/)]. [PubMed Central: [PMC6253990](https://pubmed.ncbi.nlm.nih.gov/PMC6253990/)].
26. Liu W, Wu W, Lin G, Cheng J, Zeng Y, Shi Y. Physical exercise promotes proliferation and differentiation of endogenous neural stem cells via ERK in rats with cerebral infarction. *Mol Med Rep*. 2018;**18**(2):1455–64. doi: [10.3892/mmr.2018.9147](https://doi.org/10.3892/mmr.2018.9147). [PubMed: [29901080](https://pubmed.ncbi.nlm.nih.gov/29901080/)]. [PubMed Central: [PMC6072171](https://pubmed.ncbi.nlm.nih.gov/PMC6072171/)].
27. Binte Hossain KF, Rahman MM, Sikder MT, Saito T, Hosokawa T, Kurasaki M. Inhibitory effects of selenium on cadmium-induced cytotoxicity in PC12 cells via regulating oxidative stress and apoptosis. *Food Chem Toxicol*. 2018;**114**:180–9. doi: [10.1016/j.fct.2018.02.034](https://doi.org/10.1016/j.fct.2018.02.034). [PubMed: [29454865](https://pubmed.ncbi.nlm.nih.gov/29454865/)].